



---

Year: 2019

---

## **A global database of paired leaf nitrogen and phosphorus concentrations of terrestrial plants**

Tian, Di ; Kattge, Jens ; Chen, Yahan ; Han, Wenxuan ; Luo, Yongkai ; He, Jinsheng ; Hu, Huifeng ;  
Tang, Zhiyao ; Ma, Suhui ; Yan, Zhengbing ; Lin, Quanhong ; Schmid, Bernhard ; Fang, Jingyun

**Abstract:** Nitrogen (N) and phosphorus (P) are essential components of the basic cell structure of plants. In particular, leaf N and P concentrations and their stoichiometric relationship largely determine the photosynthesis, growth, reproduction, and ecophysiological processes of plants. As important leaf functional traits, leaf N and P concentrations and their stoichiometric relationship play vital roles in indicating plant nutrient-use strategies and their evolution in terrestrial ecosystems. They also influence physiological and ecological processes in leaves (e.g., growth rate and energy metabolism) and productivity (e.g., net primary production and net ecosystem production) at ecosystem level. However, the lack of a comprehensive data set containing paired leaf N and P concentration records has distinctly limited research on nutrient stoichiometry and leaf functional traits. Here, we provide a global database of paired records of leaf N and P concentrations. A total of 11,354 individual records were acquired spanning 1,291 sites worldwide, including 201 families, 1,265 genera, and 3,227 species. The records span a latitudinal range of 45.28 °S to 68.35 °N and a longitudinal range of 155.5 °W to 168.0 °E. The variables provided for each individual record are (1) geographical location (longitude, latitude, and altitude); (2) matched leaf N and P concentrations and N:P ratio; (3) taxonomic information (family, genera, and species); (4) life form (angiosperm/gymnosperm, monocotyledonous/dicotyledonous and woody plants/herbaceous plants; note that woody plants were further divided into coniferous, deciduous broad-leaved, and evergreen broad-leaved woody species and that herbaceous plants were further divided into annual and perennial species); (5) mean annual temperature (MAT) and mean annual precipitation (MAP); and (6) soil N and P concentrations and pH value in some records. To date, this database is the world's largest database of paired leaf N and P concentrations, which contains matched information of geographical location, environmental factors, and taxa. We believe that the database will play a fundamental and crucial part of ecological stoichiometric studies. There are no copyright restrictions. When using this database, we kindly request that you cite this article, respecting all the authors' hard work during sample collection and data compilation.

DOI: <https://doi.org/10.1002/ecy.2812>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-180655>

Journal Article

Accepted Version

Originally published at:

Tian, Di; Kattge, Jens; Chen, Yahan; Han, Wenxuan; Luo, Yongkai; He, Jinsheng; Hu, Huifeng; Tang, Zhiyao; Ma, Suhui; Yan, Zhengbing; Lin, Quanhong; Schmid, Bernhard; Fang, Jingyun (2019). A

global database of paired leaf nitrogen and phosphorus concentrations of terrestrial plants. *Ecology*, 100(9):e02812.  
DOI: <https://doi.org/10.1002/ecy.2812>

**A global database of paired leaf nitrogen and phosphorus  
concentrations of terrestrial plants**

Di Tian<sup>1,2</sup>, Jens Kattge<sup>3</sup>, Yahan Chen<sup>4</sup>, Wenxuan Han<sup>5</sup>, Yongkai Luo<sup>4</sup>, Jinsheng He<sup>2</sup>,  
Huifeng Hu<sup>4</sup>, Zhiyao Tang<sup>2</sup>, Suhui Ma<sup>2</sup>, Zhengbing Yan<sup>2</sup>, Quanhong Lin<sup>1</sup>, Bernhard  
Schmid<sup>6</sup>, Jingyun Fang<sup>2\*</sup>

<sup>1</sup> College of Life Sciences, Capital Normal University, Beijing, 100048, China

<sup>2</sup> Institute of Ecology, College of Urban and Environmental Sciences, and Key  
Laboratory for Earth Surface Processes of the Ministry of Education, Peking  
University, Beijing, 100871, China

<sup>3</sup> Max-Planck-Institute for Biogeochemistry, Hans-Knöll Street 10, Jena 07745,  
Germany

<sup>4</sup> State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,  
Chinese Academy of Sciences, Beijing 100093, China

<sup>5</sup> Key Laboratory of Plant-Soil Interactions, Ministry of Education, College of  
Resources and Environmental Sciences, China Agricultural University, Beijing  
100193, China

<sup>6</sup> Department of Evolutionary Biology and Environmental Studies, University of Zurich,  
Zurich, Switzerland

**\*Corresponding author:**

Jingyun Fang, Ph.D.

Professor of Ecology

Department of Ecology

Peking University

Beijing 100871, China

Tel/Fax: +86-10-6275 6560

28 E-mail: [jyfang@urban.pku.edu.cn](mailto:jyfang@urban.pku.edu.cn)

29

**ABSTRACT.** Nitrogen (N) and phosphorus (P) are essential components of the basic cell structure of plants. In particular, leaf N and P concentrations and their stoichiometric relationship largely determine the photosynthesis, growth, reproduction and eco-physiological processes of plants. As important leaf functional traits, leaf N and P concentrations and their stoichiometric relationship play vital roles in indicating plant nutrient-use strategies and their evolution in terrestrial ecosystems. They also influence physiological and ecological processes in leaves (e.g., growth rate and energy metabolism) and at ecosystem level N or P affect productivity. However, the lack of a comprehensive dataset containing paired leaf N and P concentration records has distinctly limited research on nutrient stoichiometry and leaf functional traits. Here, we provide a global database of paired records of leaf N and P concentrations. A total of 11,354 individual records were acquired spanning 1,291 sites worldwide, including 201 families, 1,265 genera and 3,227 species. The records span a latitude range from 45.28°S to 68.35°N and a longitude range from 155.5°W to 168°E. The variables provided for each individual record are: (1) geographical location (longitude, latitude and altitude); (2) matched leaf N and P concentrations and N:P ratio; (3) taxonomic information (family, genera and species); (4) life form (angiosperm/gymnosperm, monocotyledonous/dicotyledonous and woody plants/herbaceous plants with woody plants further divided into coniferous, deciduous broad-leaved and evergreen broad-leaved woody species and herbaceous plants further divided into annual and perennial species); (5) mean annual temperature and mean annual precipitation; and (6) soil N and P concentrations and pH (partly). Up to now, this database is the world's largest database of matched leaf N and P concentrations, which contains matched information of geographical location, environmental factors, and taxa. We believe that the database will play a fundamental and crucial part in ecological stoichiometric studies. When using this database, we kindly request that you cite this article, respecting all the authors' hard work during sample collection and data compilation.

57

58 **KEY WORDS:** *leaf; nitrogen; N:P ratio; phosphorus; plant functional group;*

59 *stoichiometry; scaling exponent; terrestrial plants.*

## INTRODUCTION

At least 17 basic elements are needed for plants to grow, among which nitrogen (N) and phosphorus (P) are essential components of the basic cell structure of plants and play pivotal roles in the synthesis and transformation of protein, nucleic acids, ATP, ADP and NADP (Garten, 1976). Concentrations of N and P in plants and their balance influence not only individual plants but also the whole ecosystem, including plant growth and ecosystem productivity (Chapin et al., 1986; Sardans et al., 2012; Sullivan et al., 2014), decomposition and transformation of dead plant material, nutrient utilization and activity of symbiotic microorganisms (Treseder, 2004; Sundqvist et al., 2014), maintenance of species reproduction and diversity (Fujita et al., 2014), nutrient cycling (Asner et al., 1997), feeding by herbivores (Elser et al., 2000), energy flow and ecosystem succession (Magill et al., 2004). Especially for the leaves, the N to P stoichiometric relationship has a great influence on metabolic activities, photosynthesis and primary production of the individual plants. Therefore, leaf N and P concentrations and their stoichiometric relationships have become the focus of stoichiometry studies in plants (Sterner and Elser, 2002; Kerkhoff et al., 2006; Sardans et al., 2012).

Leaf N and P stoichiometric characteristics include N and P concentrations, N:P ratios, and N~P scaling relationships. As fundamental properties of the leaf economics spectrum (LES), leaf N and P concentrations show a strong correlation and can be used for models of relative growth rate and energy metabolism in plants (Wright et al., 2004). Empirical values of N:P ratios are often applied to indicate N or P limitation in ecosystems (Koerselman & Meuleman 1996; Güsewell, 2004). However, how does the leaf N to P stoichiometric relationship vary between regions, climates, life forms, species and other modifying variable? Understanding such variation will increase the accuracy of models and predictions about plant growth and ecosystem functioning

(Messier et al., 2016; Yan et al., 2017; He et al., 2018). Theoretically, the leaf N to P stoichiometric relationship reflects the adaptation of plants to their environment. During the process of evolution, plants developed their own mechanisms of nutrient uptake, which allows them to absorb and accumulate nutrients selectively to maintain the homeostasis of intracellular nutrient concentrations and keep a relatively constant ratio between concentrations of diverse nutrients among plant cells (Garten 1976; Sterner and Elser, 2002). Therefore, plant stoichiometric characteristics can be used as an important feature or functional trait of plant individuals and entire plant communities, highlighting different acclimations and adaptations of plants to their environment and ultimately their physiological and ecological strategies together with underlying phylogenetic processes (Kerkhoff et al., 2006; Ågren 2008; Brulheide et al., 2018).

Based on different species, communities, ecosystems and global databases, previous researchers obtained a series of widely dissimilar values of leaf N and P stoichiometric characteristics (Wright et al., 2005; McGroddy et al., 2004; Kerkhoff et al., 2006; Reich et al., 2010). This included divergent patterns of leaf N and P stoichiometric relationships along latitudinal, altitudinal and other gradients (Reich et al., 2004; Niklas et al., 2006; Zhao et al., 2016; Tian et al., 2018). It also indicated that different species, functional groups, life histories, environmental conditions (including climate factors) had complex impacts on the N to P stoichiometric relationship (Zhang et al., 2012; Yang et al., 2019). Consequently, the same plants in different regions may show significantly different N to P stoichiometric relationships, while different plant species in the same region may show similar N and P stoichiometric relationships. In order to examine such variations and the underlying driving factors, it is necessary to conduct comprehensive analyses for plants across different regions, climatic zones, taxonomic groups, life forms and other contrasts.



114 Nevertheless, further exploration of leaf N and P stoichiometric relationships have been  
115 hampered by the lack of data availability across large scales. For example, in the global  
116 database of Wright et al. (2005), only 745 records of leaf N and P (based on leaf-mass  
117 units) were available to analyze leaf N~P scaling relationships. Moreover, only  
118 countable records of N and P concentrations fell into plants from different ecosystem  
119 types including temperate forest, rainforest (Hawaii), tropical forest, tundra, wetland,  
120 alpine vegetation, and grassland. The data set of McGroddy et al. (2004) only contained  
121 three types of woody plants with 55 records located across a latitudinal range from  
122 23.5°S to 23.5°N. Niklas et al. (2005) used data from 131 vascular plants (53  
123 monocotyledonous species and 78 dicotyledonous species) to explore the relationships  
124 between leaf N to P stoichiometry and plant growth. Databases developed by Reich et  
125 al. (2004) and Reich et al. (2010) were incompletely open to the public, and the portions  
126 uploaded to the TRY database did not show matched geographical information on  
127 latitude, longitude and altitude. In summary, the completeness, uniformity and  
128 credibility of previous studies were compromised in several ways.

129

130 Considering the actual requirements and the progresses of global information about  
131 plant traits and ecosystem functioning in macroecology and biogeography (He et al.,  
132 2018), we built a comprehensive database containing an unprecedented number of  
133 records of matched leaf N and P concentrations. We combined data from the TRY-Plant  
134 Data (ref. needed here!!) and own data from large standardized field campaigns. We  
135 believe that our new database provides rich and detailed information to support research  
136 about global patterns of plant functional traits in the future. In our database we include  
137 only those records of paired N and P concentrations that come from fully developed  
138 green leaves with detailed location information. In total, 4,213 records in our database  
139 are taken from the TRY database (<https://www.try-db.org>) with specific location  
140 information; and 7,141 records come from published literature or own field data from

141 trees, shrubs and herbs in China. Duplicate records were carefully removed. Field  
142 sampling was primarily completed from July to August in the years 2000–2016. Leaf  
143 samples were dried at 60 °C for 72 h and ground in a ball mill (NM200, Retsch, Haan,  
144 Germany), and then their total N and P concentrations were measured by the Dumas  
145 combustion method using an elemental analyser (Elementar vario EL III, Elementar,  
146 Hanau, Germany; Jones, 2001). The database eventually included 11,354 records,  
147 covering 201 families, 1,265 genera, and 3,227 species. All records were derived from  
148 natural ecosystems, while ecosystems affected by fertilization or plants cultivated in  
149 the greenhouse or plantations were excluded. Life forms were divided into woody  
150 plants (coniferous, deciduous broad-leaved and evergreen broad-leaved woody species),  
151 shrubs (subshrubs), vines, herbaceous plants (annual or perennial) and ferns.

152

153 Because of standardized sampling methods, large number of records and wide spatial  
154 distribution, our database can provide comprehensive material for the study of leaf  
155 functional traits and for testing theoretical assumptions about an important ecological  
156 stoichiometric relationship, namely between leaf N and P concentrations. Furthermore,  
157 the database contains information that can help to understand nutrient cycling in  
158 ecosystems and the construction of biogeochemical models under global change.

159

## 160 **METADATA**

### 161 ***CLASS I. DATA SET DESCRIPTORS***

162 **A. Data set identity:** A global database of paired leaf nitrogen and phosphorus  
163 concentrations of terrestrial plants

164

165 **B. Data set and metadata identification codes:** Global\_Leaf\_nitrogen  
166 \_phosphorus\_concentrations.csv

167

## 168 **C. Data set description**

### 169 **1. Originators**

170 Di Tian<sup>1</sup>, Jens Kattge<sup>2</sup>, Yahan Chen<sup>3</sup>, Wenxuan Han<sup>4</sup>, Yongkai Luo<sup>3</sup>, Jinsheng He<sup>5</sup>,  
171 Huifeng Hu<sup>3</sup>, Zhiyao Tang<sup>5</sup>, Suhui Ma<sup>5</sup>, Zhengbing Yan<sup>5</sup>, Quanhong Lin<sup>1</sup>, Bernhard  
172 Schmid<sup>6</sup>, Jingyun Fang<sup>5</sup>

173 <sup>1</sup> College of Life Sciences, Capital Normal University, Beijing, 100048, China

174 <sup>2</sup> Max-Planck-Institute for Biogeochemistry, Hans-Knöll Street 10, Jena 07745,  
175 Germany

176 <sup>3</sup> State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,  
177 Chinese Academy of Sciences, Beijing 100093, China

178 <sup>4</sup> Key Laboratory of Plant-Soil Interactions, Ministry of Education, College of  
179 Resources and Environmental Sciences, China Agricultural University, Beijing 100193,  
180 China

181 <sup>5</sup> Institute of Ecology, College of Urban and Environmental Sciences, and Key  
182 Laboratory for Earth Surface Processes of the Ministry of Education, Peking University,  
183 Beijing, 100871, China

184 <sup>6</sup> Department of Evolutionary Biology and Environmental Studies, University of Zurich,  
185 Zurich, Switzerland

186

### 187 **2. Abstract**

188 Nitrogen (N) and phosphorus (P) are essential components of the basic cell structure of  
189 plants. In particular, leaf N and P concentrations and their stoichiometric relationship  
190 largely determine the photosynthesis, growth, reproduction and eco-physiological  
191 processes of plants. As important leaf functional traits, leaf N and P concentrations and  
192 their stoichiometric relationship play vital roles in indicating plant nutrient-use  
193 strategies and their evolution in terrestrial ecosystems. They also influence  
194 physiological and ecological processes in leaves (e.g., growth rate and energy

metabolism) and at ecosystem level N or P affect productivity. However, the lack of a comprehensive dataset containing matched leaf N and P concentration records has distinctly limited research on nutrient stoichiometry and leaf functional traits. Here, we provide a global database of paired records of leaf N and P concentrations. A total of 11,354 individual records were acquired spanning 1,291 sites worldwide, including 201 families, 1,265 genera and 3,227 species. The records span a latitude range from 45.28°S to 68.35°N and a longitude range from 155.5°W to 168°E. The variables provided for each individual record are: (1) geographical location (longitude, latitude and altitude); (2) matched leaf N and P concentrations and N:P ratio; (3) taxonomic information (family, genera and species); (4) life form (angiosperm/gymnosperm, monocotyledonous/dicotyledonous and woody plants/herbaceous plants with woody plants further divided into coniferous, deciduous broad-leaved and evergreen broad-leaved woody species and herbaceous plants further divided into annual and perennial species); (5) mean annual temperature and mean annual precipitation; and (6) soil N and P concentrations and pH (partly). Up to now, this database is the world's largest database of matched leaf N and P concentrations, which contains matched information of geographical locations, environmental factors, and taxa. We believe that the database will play a fundamental and crucial part in ecological stoichiometric studies. When using this database, we kindly request that you cite this article, respecting all the authors' hard work during sample collection and data compilation.

**3. Key words:** *leaf, nitrogen, N:P ratio, phosphorus, plant functional group, stoichiometry, scaling exponent, terrestrial plants*

## **CLASS II. RESEARCH ORIGIN DESCRIPTORS**

### **A. Overall project description**

**1. Identity:** A global database of paired leaf nitrogen and phosphorus concentrations of

222 terrestrial plants

223 **2. Originators:** J.Y. Fang, D. Tian, J. Kattge, W.X. Han, Y.H. Chen, Y.K. Luo, H.F. Hu,  
224 Z.Y. Tang, J.S. He, S.H. Ma, Z.B. Yan, Q.H. Lin, B. Schmid

225 **3. Period of study:** Field sampling was conducted during 2000–2016. The database  
226 was compiled between 2015 and 2018.

227 **4. Objectives:** (i) To document global variation in leaf nitrogen and phosphorus  
228 concentrations of terrestrial plants and examine leaf N and P stoichiometric  
229 relationships among different life forms, phylogenetic taxa, biomes, plant communities  
230 and sites; and (ii) to provide a broad-scale comprehensive dataset for verifying  
231 important leaf functional traits and classic theories in plant ecological stoichiometry  
232 across individual, species, genus, family, community, ecological biomes and global  
233 scales.

234 **5. Sources of funding:** This work was supported by the National Natural Science  
235 Foundation of China (grant no. 31800397, 31321061 and 31330012), National Key  
236 R&D Program of China (grant no. 2017YFC0503900) and the TRY initiative on plant  
237 traits (<http://www.try-db.org>). The TRY database is hosted at the Max Planck Institute  
238 for Biogeochemistry (Jena, Germany) and supported by DIVERSITAS/Future Earth,  
239 the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig and  
240 EU project BACI [grant ID 640176].

241

## 242 **B. Specific subproject description**

### 243 **1. Data sources**

244 The dataset was partly compiled from an open data source (i.e. the TRY-Plant Trait  
245 Database, <https://www.try-db.org/TryWeb/Data.php>; Kattge et al., 2011), an integrated  
246 data set of China established by Han et al. (2005) and four major original sampling  
247 studies (He et al., 2008; Chen et al., 2013; Yang et al., 2016; Luo, 2017 *unpublished*).

248

We collected only those paired records of N and P concentrations in green leaves with detailed location information and excluded other records without site information or with unpaired N-P records. Specifically, the sub-datasets in TRY-Plant Trait Database were carefully compiled: Abisko & Sheffield Database (ID1, Quested et al., 2003), GLOPNET-Global Plant Trait Network Database (ID20, Wright et al., 2004), The RAINFOR Plant Trait Database (ID34, Fyllas et al., 2009), Sheffield Database (ID37, Cornelissen et al., 2003), The VISTA Plant Trait Database (ID45, Garnier et al., 2007), Leaf and Whole Plant Traits Database (ID50, Kazakou et al., 2006), Tropical Plant Traits From Borneo Database (ID51, Swaine, 2007), Wetland Dunes Database (ID56, Bakker et al., 2005), Global Leaf Robustness and Physiology Database (ID87, Niinemets, 2001), The Netherlands Plant Traits Database (ID88, Ordonez et al., 2010), Ukraine Wetlands Plant Traits Database (ID90, van Bodegom, *unpublished*), Global A, N, P, SLA Database (ID94, Reich et al., 2009), Traits from Subarctic Plant Species Database (ID105, Freschet et al., 2010), Global <sup>15</sup>N Database (ID130, Craine et al., 2005), Functional Traits of Graminoids in Semi-Arid Steppes Database (ID152, Adler et al., 2004), Traits of Hemiparasitic Plants (ID166, Demey et al., 2013) and Plant Traits for Pinus and Juniperus Forests in Arizona (ID193, Laughlin et al., 2011).

In Han et al. (2005), 2,094 records of paired leaf N and P concentrations from 127 sampling sites across China were collected from published literature. Among these records, 1,979 were incorporated into the current database because they contained detailed taxonomic information about family, genus and species. The other data from four major original studies (He et al., 2008; Chen et al., 2013; Yang et al., 2016; Luo, 2017 *unpublished*) were sampled using standardized collection protocols in the summer (see details in **Experimental Design and Research Method**).

## **2. Site description**

276 **a. Sites and Geography**

277 Sites include natural forest, grassland, shrubland and wetland ecosystems across the  
278 globe. The terrestrial sites include the following continents: Asia, Africa, Europe,  
279 Oceania (including Hawaii) and America (North America, Central America and South  
280 America). In total, coverage includes boreal, temperate and tropical ecosystems except  
281 Antarctica.

282 **b. Habitat**

283 These sites cover a wide array of natural habitats from humid tropical forests to boreal  
284 forests, arid, semi-arid grasslands, shrublands and alpine ecosystems. The typical  
285 vegetation types include desert, scrub, steppe, meadow, shrublands, boreal forests,  
286 temperate deciduous forests, subtropical and tropical evergreen forests, as well as  
287 wetlands.

288 **c. Geology**

289 Sites include terrestrial sites scattered across the globe and therefore present various  
290 geological formations. For example, data from Hawaii cover old and young site ages  
291 and thus soil developmental stages along a chronosequence.

292 **d. Site history**

293 All sites are in natural ecosystems without or with weak anthropogenic influences.  
294 Greenhouse cultivations and artificial plantations are not included in the database.

295 **e. Climate**

296 Climate ranges from boreal, temperate to tropical regions with substantial annual  
297 variations in both temperature and rainfall. For example, mean annual temperature  
298 ranges across observations from  $-11.4$  to  $27.8$  °C and mean annual precipitation ranges  
299 from 17 to 3922 mm.

300

301 **3. Experimental design**

302 Leaf samples from forests (Chen et al., 2013), shrublands (Yang et al., 2016; Luo, 2017)

303 and grasslands (He et al., 2006, 2008; Geng et al., 2017) across China were collected  
304 in summer from 2000 to 2016. In these natural ecosystems, plots were selected  
305 according to the vegetation of local sites and then at least 5 fully expanded sun leaves  
306 of plant individuals of each species in the plots were collected and oven-dried at 60 °C  
307 in the laboratory. After being grounded in ball mills (NM200, Retsch, Haan, Germany),  
308 the powder of leaf samples was used to measure leaf N concentrations with an elemental  
309 analyzer (2400 II CHNS/O Elemental Analyzer, Perkin-Elmer, USA) with a  
310 combustion temperature of 950 °C and a reduction temperature of 640 °C. Leaf P  
311 concentrations were determined with the molybdate/ascorbic acid method after H<sub>2</sub>SO<sub>4</sub>-  
312 HClO<sub>4</sub> digestion (Jones, 2001). The absorbance of each sample for P concentration  
313 detection was measured at 700 nm after 20 min of adding molybdenum-stibium-  
314 ascorbic acid reagent (Jones, 2001).

315

316 Soil samples at 0–10 cm depth in forests were taken in plots where the leaf samples  
317 were sampled. Three to five soil cores in each plot were randomly collected and mixed  
318 evenly. Then soil total N and P concentrations were detected through air-drying, sieving  
319 and elemental analyzer similar to the measurement processes of leaf N and P  
320 concentrations. The average values of soil N and P concentrations in three plots at the  
321 same site were used to represent the soil N and P concentrations of local plant species  
322 (Chen et al., 2013). In shrublands and grasslands, subsamples from three pits at 0–10  
323 cm depth were mixed and detected. Soil pH was measured in water suspension with a  
324 1:2.5 soil:water ratio using a benchtop pH meter (SevenEasy Plus, Mettler Toledo,  
325 Switzerland) or in both 0.01 M CaCl<sub>2</sub> and bi-distilled H<sub>2</sub>O potentiometrically (Geng et  
326 al., 2006). The average values of soil variables from three soil subsamples near the plant  
327 samples were used to represent their corresponding soil N and P concentrations.

328

#### 329 **4. Research methods**



330 The data processing involved four steps: (1) standardization of leaf N and P records and  
331 supplementary variables including matched soil N and P concentrations, soil pH and  
332 brief descriptions of their locations, (2) assignment of a consensus of family, genus and  
333 species names, (3) standardization of growth-form classification to each species, and  
334 (4) extraction of mean annual temperature (MAT) and mean annual precipitation (MAP).

335

336 For the first step, paired N and P concentrations of green leaves with detailed location  
337 information (latitude and longitude) were incorporated into the database and other  
338 records without site information or with unpaired N-P records were excluded.  
339 According to the related references with geographic information on latitude, longitude  
340 and altitude (for some records), we located the sites of each individual and checked  
341 their local vegetation types and biomes.

342

343 Then, we checked and matched the species names of each individual with standardized  
344 abbreviations removing special characters and numbers in the names. Furthermore, we  
345 used the following literature to identify plant growth forms and phylogenetic taxonomy:  
346 the Flora of China (<http://frps.eflora.cn/>), Angiosperm Phylogeny Website  
347 (<http://www.mobot.org/MOBOT/research/APweb/>) (Stevens, 2001), Useful Tropical  
348 Plants (<http://tropical.theferns.info/>), Wikipedia (<https://en.wikipedia.org/wiki>) and  
349 Australian Native Plants (<https://www.anbg.gov.au/index.html>). To avoid  
350 inconsistencies in plant names and classifications, the taxonomy of all records was  
351 homogenized following the Flora of China (<http://frps.eflora.cn/>). A plant growth form  
352 dataset for the New World established by Engelmann et al. (2016) was also used in our  
353 dataset to determine plant growth forms. All plants were assigned to one of six growth  
354 forms: ferns, herbs, subshrubs (e.g. *Artemisia* species), shrubs, trees or vines.  
355 Subsequently, the six growth forms except for subshrubs were further divided into  
356 deciduous coniferous species, deciduous broadleaved species, evergreen coniferous

species, evergreen broadleaved species, dicotyledonous herbs, monocotyledonous herbs and semi-deciduous species on the basis of their morphology and physiology. Here, a small part of species with fleshy leaves, thorns and other hardly interpretable leaf types were only grouped into evergreen or deciduous species.

Finally, the climatic factors including MAT and MAP were predominately compiled from the original studies. The other MAT/MAP variables not provided in the original publications were estimated through data extraction from a global climatic data set (<http://worldclim.org/version2>) with a resolution of  $0.0083 \times 0.0083$  (*ca.* 1 km<sup>2</sup>). Additionally, the corresponding data of soil total phosphorus density in the top 50 cm soil were extracted from Global Gridded Soil Phosphorus Distribution Maps (Yang et al., 2014, <http://dx.doi.org/10.3334/ORNLDAAAC/1223>) for those compiled leaf N and P concentration records without soil nutrient content information.

## **5. Project personnel**

Many people including graduate students from Peking University, workers and researchers have contributed to the development of this project, helping to take the measurements in field sampling, investigation and data compiling.

## ***CLASS III. DATA SET STATUS AND ACCESSIBILITY***

### **A. Status**

**1. Latest update:** May 2019.

**2. Metadata status:** Metadata are complete.

**3. Storage location and medium:** Parts of the data set are stored in raw data order at the TRY Data Portal (ID53 and ID181, contributed by Jingyun Fang and Wenxuan Han) hosted at <https://www.try-db.org/de/DatasetDetails.php>. The whole dataset will be stored as Supporting Information to this publication in the Ecological Society of

America's journal *Ecology*. The dataset is also available via direct access to the contact person. The dataset is also stored in non-proprietary digital form by the contact person. Electronic or paper copies of data sources (articles, reports, theses, etc.) are also available via direct access to the corresponding persons.

## **B. Data verification**

The dataset has passed through several rounds of review as the data contributors have published several studies based on parts of the dataset (e.g. Han et al., 2005; Chen et al., 2013; Geng et al., 2016; Yang et al., 2016; Yan et al., 2017; Tian et al., 2018), which ensured the accuracy of the units of variables and the matching of leaf N and P concentrations and the associated variables. Moreover, the data compilers have also gone through extensive data verification by checking credible ranges and outliers of all variables in the database.

## **C. Accessibility**

### **1. Contact person:**

Jingyun Fang, email: [jyfang@urban.pku.edu.cn](mailto:jyfang@urban.pku.edu.cn); Tel/Fax: 86-10-6275 6560; Affiliation: Institute of Ecology, College of Urban and Environmental Sciences, Peking University, 5 Yiheyuan Rd., Beijing 100871, China.

Di Tian, email: [tiandi@cnu.edu.cn](mailto:tiandi@cnu.edu.cn); Tel/Fax: 86-10-6890 2374; Affiliation: College of Life Sciences, Capital Normal University, 105 West Three Ring North Rd., Beijing, 100048, China

**2. Copyright restrictions:** No copyright restrictions.

**3. Proprietary restrictions:** None. When using the dataset, we request the users to cite this paper, recognizing their use of the data in publications, research proposals, websites and other outlets.

**4. Costs:** None.

411

412 **CLASS IV: DATA STRUCTURAL DESCRIPTORS**

413 **A. Data set file**

414 **1. Identity:** Leaf\_Nitrogen\_and\_phosphorus\_concentrations\_data.csv.

415 **2. Size:** 2.11 MB.

416 **3. Format:** comma-delimited plain text, no compression.

417 **4. Header information:** The header row indicates variable names as described in Table  
418 1 below. The dash means that the field is not applicable to that dataset.

419

420 **Table 1.** Definitions of the variables included in the data set file  
421 “Leaf\_Nitrogen\_and\_phosphorus\_concentrations\_data.csv”. The dash means that the  
422 field is not applicable to that dataset.

Column	Variable	Variable description	Unit	Range
1	Individual_ID	Unique code identifying each individual	-	1 ~ 11355
2	Leaf_N	Leaf N concentration	mg g <sup>-1</sup>	2.48 ~ 66.80
3	Leaf_P	Leaf P concentration	mg g <sup>-1</sup>	0.08 ~ 9.59
4	N_P_Ratio	Leaf N:P ratio	-	1.63 ~ 99.23
5	Latitude	The original latitude: a negative value indicates southern latitude, while a positive value represents northern latitude	°	-45.28 ~ 68.35
6	Longitude	The original longitude: a negative value indicates western longitude, while a positive value represents eastern longitude	°	-155.5 ~ 168.0
7	Altitude	Altitude	m	-2 ~ 5249

8	Location_description	A brief description about the location of the record	-	-
9	MAT	Mean annual temperature	°C	-11.4 ~ 27.8
10	MAP	Mean annual precipitation	mm	17 ~ 3922
11	Soil_N	Soil total N concentration	mg g <sup>-1</sup>	0.118 ~ 19.53
12	Soil_P	Soil total P concentration	mg g <sup>-1</sup>	0.022 ~ 3.981
13	Soil_P_density	Soil total phosphorus density in the top 50 cm soil extracted from Global Gridded Soil Phosphorus Distribution Maps	g m <sup>-2</sup>	69.00 ~ 1323.81
14	Soil_pH	Soil pH	-	3.3 ~ 8.9
15	Phylum	Phylum including angiosperm, gymnosperm and fern	-	-
16	Cotyledon	Angiosperms divided into dicot, monocot and fern	-	-
17	Family_New	Family names according to Angiosperm Phylogeny Website ( <a href="http://www.mobot.org/MOBOT/research/APweb/">http://www.mobot.org/MOBOT/research/APweb/</a> ) (Stevens, 2001)	-	-
18	Family	Family names according to Flora of China ( <a href="http://frps.eflora.cn/">http://frps.eflora.cn/</a> )	-	-
19	Genus	Genus name determined through taxonomic checking	-	-

20	Species	Species name determined through taxonomic checking	-	-
21	Life_form1	Assignment to life form on the basis of morphology and physiology, including ferns (Fern), herbs (H), subshrubs (Subs), shrubs (S), trees (T) and vines (V).	-	-
22	Life_form2	Assignment to life form including deciduous coniferous species (Dc), deciduous broadleaved species (Db), evergreen coniferous species (Ec), evergreen broadleaved species (Eb), dicotyledonous herbs (Hf), monocotyledonous herbs (Hg), semi-deciduous species (Semi). A small part of species with fleshy leaves, thorns and other hardly interpretable leaf types were only distinguished to evergreen (E) or deciduous (D).	-	-
23	Leguminous	Legume (Y) or not (N)	-	-
24	dataset_name	Data sources	-	-

423

424 **B. Data anomalies**

425 NA in the database indicates missing data or no matching information. Some records

426 lack the corresponding data of altitude, soil total N and P concentrations and soil pH.

427

## 428 ***CLASS V: SUPPLEMENTAL DESCRIPTORS***

### 429 **A. Data acquisition**

#### 430 **1. Data forms or acquisition methods**

431 The dataset was compiled from an open data source (i.e. the TRY-Plant Trait Database,  
432 <https://www.try-db.org/TryWeb/Data.php>; Kattge et al. 2011), an integrated data set of  
433 China established by Han et al. (2005) and four major original sampling studies (He et  
434 al., 2008; Chen et al., 2013; Yang et al., 2016; Luo, 2017 *unpublished*) (see section  
435 II.B.1). All the records from these sub-datasets were finally integrated into one Excel  
436 sheet.

#### 437 **2. Location of completed data forms**

438 Some original sub-databases of our dataset can be found in the TRY-Plant Trait  
439 Database (<https://www.try-db.org/TryWeb/Data.php>) using Data Explorer of “ID”. The  
440 completed data forms can be found through the contact person.

#### 441 **3. Data entry verification procedures**

442 We removed duplicated records from the data set by eye.

443

### 444 **B. Quality control procedures**

445 We collected only those paired records of N and P concentrations in green leaves with  
446 detailed location information and excluded other records without site information or  
447 with unpaired N-P records. In the dataset, all the plants were distinguished to six growth  
448 forms including ferns, herbs, subshrubs, shrubs, trees and vines. A small number of  
449 species with fleshy leaves, thorns and other leaf morphology that was hard to classify  
450 into a leaf type, they were only grouped into evergreen or deciduous species without  
451 further division. The taxonomy of all records was homogenized following the Flora of  
452 China (<http://frps.eflora.cn/>).

453

#### 454 **C. Related materials**

455 The references for data sources from TRY-Plant Trait Database ([https://www.try-](https://www.try-db.org/TryWeb/Data.php)  
456 [db.org/TryWeb/Data.php](https://www.try-db.org/TryWeb/Data.php)) have been introduced in section II.B.1 and listed in the  
457 section “Literature Cited”.

458

#### 459 **E. Archiving**

460 See the data sources in section II.B.1. The data processing was introduced in section  
461 II.B.4.

462

#### 463 **F. Publication and results**

464 To better explore and interpret nutrient stoichiometry of terrestrial plants, Peking  
465 University led our field investigations and sampling across China during the past two  
466 decades. Our colleagues have published the following publications based on parts of  
467 the current data set:

468 Chen, Y. H., W. X. Han, L. Y. Tang, Z. Y. Tang, and J. Y. Fang. 2013. Leaf nitrogen and  
469 phosphorus concentrations of woody plants differ in responses to climate, soil and  
470 plant growth form. *Ecography* 36: 178-184.

471 Geng, Y., Z. H. Wang, C. Z. Liang, J. Y. Fang, F. Baumann, P. Kühn, T. Scholten, and  
472 J.-S. He. 2012. Effect of geographical range size on plant functional traits and the  
473 relationships between plant, soil and climate in Chinese grasslands. *Global*  
474 *Ecology and Biogeography* 21: 416-427.

475 Geng, Y., W. H. Ma, L. Wang, F. Baumann, P. Kühn, T. Scholten, and J.-S. He. 2017.  
476 Linking above- and belowground traits to soil and climate variables: an integrated  
477 database on China's grassland species. *Ecology* 98: 1471.

478 Guo, Y. P., X. Yang, C. Schöb, Y. X. Jiang, and Z. Y. Tang. 2017. Legume Shrubs Are  
479 More Nitrogen-Homeostatic than Non-legume Shrubs. *Frontiers in Plant Science*



480 8: 1662.

481 Han, W. X., J. Y. Fang, D. L. Guo, and Y. Zhang. 2005. Leaf nitrogen and phosphorus  
 482 stoichiometry across 753 terrestrial plant species in China. *New Phytologist* 168:  
 483 377-385.

484 Han, W. X., J. Y. Fang, P. B. Reich, F. Ian Woodward, and Z. H. Wang. 2011.  
 485 Biogeography and variability of eleven mineral elements in plant leaves across  
 486 gradients of climate, soil and plant functional type in China. *Ecology Letters* 14:  
 487 788-796.

488 He, J.-S., J. Y. Fang, Z. H. Wang, D. L. Guo, D. F. B. Flynn, and Z. Geng. 2006.  
 489 Stoichiometry and large-scale patterns of leaf carbon and nitrogen in the grassland  
 490 biomes of China. *Oecologia* 149: 115-122.

491 He, J.-S., L. Wang, D. F. B. Flynn, X. P. Wang, W. H. Ma, and J. Y. Fang. 2008. Leaf  
 492 nitrogen: phosphorus stoichiometry across Chinese grassland biomes. *Oecologia*  
 493 155: 301-310.

494 He, J.-S., X. P. Wang, D. F. B. Flynn, L. Wang, B. Schmid, and J. Y. Fang. 2009.  
 495 Taxonomic, phylogenetic, and environmental trade-offs between leaf productivity  
 496 and persistence. *Ecology* 90: 2779-2791.

497 He, J.-S., X. P. Wang, B. Schmid, D. F. B. Flynn, X. F. Li, P. B. Reich, and J. Y. Fang.  
 498 2010. Taxonomic identity, phylogeny, climate and soil fertility as drivers of leaf  
 499 traits across Chinese grassland biomes. *Journal of Plant Research* 123: 551-561.

500 He, J.-S., Z. H. Wang, X. P. Wang, B. Schmid, W. Y. Zuo, M. Zhou, C. Y. Zheng, M. F.  
 501 Wang, and J. Y. Fang. 2006b. A test of the generality of leaf trait relationships on  
 502 the Tibetan Plateau. *New Phytologist* 170: 835-848.

503 Tang, Z. Y., W. T. Xu, G. Y. Zhou, Y. F. Bai, J. X. Li, X. L. Tang, D. M. Chen, Q. Liu,  
 504 W. H. Ma, G. M. Xiong, H. L. He, N. P. He, Y. P. Guo, Q. Guo, J. L. Zhu, W. X.  
 505 Han, H. F. Hu, J. Y. Fang, and Z. Q. Xie. 2018. Patterns of plant carbon, nitrogen,  
 506 and phosphorus concentration in relation to productivity in China's terrestrial

ecosystems. *Proceedings of the National Academy of Sciences* 115: 4033-4038.

Tian, D., Z. B. Yan, K. J. Niklas, W. X. Han, J. Kattge, P. B. Reich, Y. K. Luo, Y. H. Chen, Z. Y. Tang, H. F. Hu, I. J. Wright, B. Schmid, and J. Y. Fang. 2018. Global leaf nitrogen and phosphorus stoichiometry and their scaling exponent. *National Science Review* 5: 728-739.

Yan, Z. B., P. Li, Y. H. Chen, W. X. Han, and J. Y. Fang. 2016. Nutrient allocation strategies of woody plants: an approach from the scaling of nitrogen and phosphorus between twig stems and leaves. *Scientific Reports* 6.

Yan, Z. B., D. Tian, W. X. Han, Z. Y. Tang, and J. Y. Fang. 2017. An assessment on the uncertainty of the nitrogen to phosphorus ratio as a threshold for nutrient limitation in plants. *Ann Bot* 120: 937.

Yang, X., X. L. Chi, C. J. Ji, H. Y. Liu, W. H. Ma, A. Mohhammat, Z. Y. Shi, X. P. Wang, S. L. Yu, M. Yue, and Z. Y. Tang. 2016. Variations of leaf N and P concentrations in shrubland biomes across northern China: phylogeny, climate and soil. *Biogeosciences* 13: 4429-4438.

522

## 523 **G. History of data set usage**

- 524 1. Data request history. None.
- 525 2. Data set update history. None.
- 526 3. Review history. None.
- 527 4. Questions and comments from secondary users. None

528

## 529 **ACKNOWLEDGEMENTS**

530 More than 200 investigators joined us during the long-term continual field sampling  
 531 and data collection across China. We sincerely appreciate their kind support. We also  
 532 thank Wenjing Fang, Ming Ouyang, Aijun Xing, Mengying Zhao, Hanyue Huang,  
 533 Anwar Eziz, Chao Li and Chenjun Ji for their help during our data compiling. This

534 research was supported by the National Natural Science Foundation of China (grant no.  
535 31800397, 31321061 and 31330012), National Key R&D Program of China (grant no.  
536 2017YFC0503900) and the TRY initiative on plant traits (<http://www.try-db.org>). The  
537 TRY database is hosted at the Max Planck Institute for Biogeochemistry (Jena,  
538 Germany) and supported by DIVERSITAS/Future Earth, the German Centre for  
539 Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig and EU project BACI  
540 [grant ID 640176].

541

#### 542 **LITERATURE CITED:**

543 Adler, P. B., D. G. Milchunas, W. K. Lauenroth, O. E. Sala, and I. C. Burke. 2004.  
544 Functional traits of graminoids in semi-arid steppes: a test of grazing histories.  
545 Journal of Applied Ecology 41: 653-663.

546 Aerts, R. 1996. Nutrient resorption from senescing leaves of perennials: are there  
547 general patterns? Journal of Ecology: 597-608.

548 Ågren, G. I., E. Bosatta, and A. H. Magill. 2001. Combining theory and experiment to  
549 understand effects of inorganic nitrogen on litter decomposition. Oecologia 128:  
550 94-98.

551 Ågren, G. I. 2008. Stoichiometry and nutrition of plant growth in natural communities.  
552 Annual Review of Ecology Evolution & Systematics 39: 153-170.

553 Asner, G. P., T. R. Seastedt, and A. R. Townsend. 1997. The decoupling of terrestrial  
554 carbon and nitrogen cycles. BioScience 47: 226-234.

555 Bakker, C., J. Rodenburg, and P. M. van Bodegom. 2005. Effects of Ca- and Fe-rich  
556 seepage on P availability and plant performance in calcareous dune soils. Plant  
557 Soil 275: 111-122.

558 Bauer, G. A., F. A. Bazzaz, R. Minocha, S. Long, A. Magill, J. Aber, and G. M. Berntson.  
559 2004. Effects of chronic N additions on tissue chemistry, photosynthetic capacity,  
560 and carbon sequestration potential of a red pine (*Pinus resinosa* Ait.) stand in the

561 NE United States. *Forest Ecology and Management* 196: 173-186.

562 Bruelheide, H., J. Dengler, O. Purschke, J. Lenoir, B. Jiménez-Alfaro, S. M. Hennekens,  
563 Z. Botta-Dukát, M. Chytrý, R. Field, F. Jansen , J. Kattge, V.D. Pillar, F. Schrodte,  
564 M. D. Mahecha, R. K. Peet, B. Sandel, P. van Bodegom, J. Altman, E. Alvarez-  
565 Dávila, M. A. S. Arfin Khan, F. Attorre, I. Aubin, C. Baraloto, J. G. Barroso,  
566 M. Bauters, E. Bergmeier, I. Biurrun, A. D. Bjorkman, B. Blonder, A. Čarni,  
567 L. Cayuela, T. Černý, J. H. C. Cornelissen, D. Craven, M. Dainese, G. Derroire,  
568 M. D. Sanctis, S. Díaz, J. Doleža, W. Farfan-Rios, T. R. Feldpausch, N. J. Fenton,  
569 E. Garnier, G. R. Guerin, A. G. Gutiérrez, S. Haider, T. Hattab, G. Henry,  
570 B. Hérault, P. Higuchi, N. Hölzel, J. Homeier, A. Jentsch , N. Jürgens, Z. Kacki,  
571 D. N. Karger, M. Kessler, M. Kleyer, I. Knollová, A. Y. Korolyuk, I. Kühn,  
572 D. C. Laughlin, F. Lens, J. Loos, F. Louault, M. I. Lyubenova, Y. Malhi,  
573 C. Marcenò, M. Mencuccini, J. V. Müller, J. Munzinger, I. H. Myers-Smith,  
574 D. A. Neill, Ü. Niinemets, K. H. Orwin, W. A. Ozinga, J. Penuelas, A. Pérez-  
575 Haase , P. Petřík, O. L. Phillips, M. Pärtel, P. B. Reich, C. Römermann,  
576 A. V. Rodrigues, F. M. Sabatini, J. Sardans, M. Schmidt, G. Seidler, J. E. S.  
577 Espejo, M. Silveira, A. Smyth, M. Sporbert, J.-C. Svenning, Z.Y. Tang,  
578 R. Thomas, I. Tsiripidis, K. Vassilev, C. Violle, R. Virtanen, E. Weiher, E. Welk ,  
579 K. Wesche, M. Winter, C. Wirth, and U. Jandt. 2018. Global trait–environment  
580 relationships of plant communities. *Nature ecology & evolution* 2: 1906.

581 Chapin III, F. S., G. R. Shaver, and R. A. Kedrowski. 1986. Environmental controls  
582 over carbon, nitrogen and phosphorus fractions in *Eriophorum vaginatum* in  
583 Alaskan Tussock Tundra. *Journal of Ecology* 74: 167-195.

584 Chen, Y. H., W. X. Han, L. Y. Tang, Z. Y. Tang, and J. Y. Fang. 2013. Leaf nitrogen and  
585 phosphorus concentrations of woody plants differ in responses to climate, soil and  
586 plant growth form. *Ecography* 36: 178-184.

587 Cornelissen, J. H. C., B. Cerabolini, P. Castro-Díez, P. Villar-Salvador, G. Montserrat-

588 Martı́, J. P. Puyravaud, M. Maesto, M. J. A. Werger, and R. Aerts. 2003. Functional  
589 traits of woody plants: correspondence of species rankings between field adults  
590 and laboratory-grown seedlings? *Journal of Vegetation Science* 14: 311-322.

591 Craine, J. M., W. G. Lee, W. J. Bond, R. J. Williams, and L. C. Johnson. 2005.  
592 Environmental constraints on a global relationship among leaf and root traits of  
593 grasses. *Ecology* 86: 12-19.

594 Demey, A., J. Staelens, L. Baeten, P. Boeckx, M. Hermy, J. Kattge, and K. Verheyen.  
595 2013. Nutrient input from hemiparasitic litter favors plant species with a fast-  
596 growth strategy. *Plant Soil* 371: 53-66.

597 Elser, J. J., W. J. O'Brien, D. R. Dobberfuhl, and T. E. Dowling. 2000. The evolution  
598 of ecosystem processes: growth rate and elemental stoichiometry of a key  
599 herbivore in temperate and arctic habitats. *Journal of Evolutionary Biology* 13:  
600 845-853.

601 Engemann, K., B. Sandel, B. Boyle, B. J. Enquist, P. M. Järgensen, J. Kattge, B. J.  
602 McGill, N. Morueta-Holme, R. K. Peet, N. J. Spencer, C. Violle, S. K. Wiser, and  
603 J.-C. Svenning. 2016. A plant growth form dataset for the New World. *Ecology* 97:  
604 3243.

605 Fyllas, N. M., S. Patiño, T. R. Baker, G. B. Nardoto, L. A. Martinelli, C. A. Quesada,  
606 R. Paiva, M. Schwarz, V. Horna, L. M. Mercado, A. Santos, L. Arroyo, E. M.  
607 Jimenez, F. J. Luizão, D. A. Neill, N. Silva, A. Prieto, A. Rudas, M. Silviera, I. C.  
608 G. Vieira, G. Lopez-Gonzalez, Y. Malhi, O. L. Phillips, and J. Lloyd. 2009. Basin-  
609 wide variations in foliar properties of Amazonian forest: phylogeny, soils and  
610 climate. *Biogeosciences* 6: 2677-2708.

611 Fujita, Y., H. O. Venterink, P. M. van Bodegom, J. C. Douma, G. W. Heil, N. Hölzel, E.  
612 Jabłońska, W. Kotowski, T. Okruszko, P. Pawlikowski, P. C. de Ruiter, and M. J.  
613 Wassen. 2014. Low investment in sexual reproduction threatens plants adapted to  
614 phosphorus limitation. *Nature* 505: 82.

615 Garnier, E., S. Lavorel, P. Ansquer, H. Castro, P. Cruz, J. Dolezal, O. Eriksson, C.  
 616 Fortunel, H. Freitas, C. Golodets, K. Grigulis, C. Jouany, E. Kazakou, J. Kigel, M.  
 617 Kleyer, V. Lehsten, J. Lepš, T. Meier, R. Pakeman, M. Papadimitriou, V. P.  
 618 Papanastasis, H. Quested, F. Quétier, M. Robson, C. Roumet, G. Rusch, C. Skarpe,  
 619 M. Sternberg, J. P. Theau, A. Thébault, D. Vile, and M. P. Zarovali. 2007.  
 620 Assessing the effects of land-use change on plant traits, communities and  
 621 ecosystem functioning in grasslands: a standardized methodology and lessons  
 622 from an application to 11 European sites. *Annals of Botany* 99: 967-985.

623 Garten, C. T. 1976. Correlations between concentrations of elements in plants. *Nature*  
 624 261: 686-688.

625 Geng, Y., W. H. Ma, L. Wang, F. Baumann, P. Kühn, T. Scholten, and J.-S. He. 2017.  
 626 Linking above- and belowground traits to soil and climate variables: an integrated  
 627 database on China's grassland species. *Ecology* 98: 1471.

628 Han, W. X., J. Y. Fang, D. L. Guo, and Y. Zhang. 2005. Leaf nitrogen and phosphorus  
 629 stoichiometry across 753 terrestrial plant species in China. *New Phytologist* 168:  
 630 377-385.

631 He, J.-S., J. Y. Fang, Z. H. Wang, D. L. Guo, D. F. B. Flynn, and Z. Geng. 2006.  
 632 Stoichiometry and large-scale patterns of leaf carbon and nitrogen in the grassland  
 633 biomes of China. *Oecologia* 149: 115-122.

634 He, J.-S., L. Wang, D. F. B. Flynn, X. P. Wang, W. H. Ma, and J. Y. Fang. 2008. Leaf  
 635 nitrogen: phosphorus stoichiometry across Chinese grassland biomes. *Oecologia*  
 636 155: 301-310.

637 Jones, J. B. Laboratory Guide for Conducting Soil Tests and Plant Analysis, CRC Press,  
 638 New York, NY, 2001.

639 Kattge, J., S. Diaz, S. Lavorel, I. C. Prentice, P. Leadley, G. Bönsch, E. Garnier, M.  
 640 Westoby, P. B. Reich, I. J. Wright, J. H. C. Cornelissen, C. Violle, S. P. Harrison,  
 641 P. M. Van Bodegom, M. Reichstein, B. J. Enquist, N. A. Soudzilovskaia, D. D.

642 Ackerly, M. Anand, O. Atkin, M. Bahn, T. R. Baker, D. Baldocchi, R. Bekker, C.  
 643 C. Blanco, B. Blonder, W. J. Bond, R. Bradstock, D. E. Bunker, F. Casanoves, J.  
 644 Cavender-Bares, J. Q. Chambers, F. S. Chapin III, J. Chave, D. Coomes, W. K.  
 645 Cornwell, J. M. Craine, B. H. Dobrin, L. Duarte, W. Durka, J. Elser, G. Esser, M.  
 646 Estiarte, W. F. Fagan, J. FANG, F. Fernández-Méndez, A. Fidelis, B. Finegan, O.  
 647 Flores, H. Ford, D. Frank, G. T. Freschet, N. M. Fyllas, R. V. Gallagher, W. A.  
 648 Green, A. G. Gutierrez, T. Hickler, S. I. Higgins, J. G. Hodgson, A. Jalili, S. Jansen,  
 649 C. A. Joly, A. J. Kerkhoff, D. Kirkup, K. Kitajima, M. Kleyer, S. Klotz, J. M. H.  
 650 Knops, K. Kramer, I. Kühn, H. Kurokawa, D. Laughlin, T. D. Lee, M. Leishman,  
 651 F. Lens, T. Lenz, S. L. Lewis, J. Lloyd, J. Llusià, F. Louault, S. Ma, M. D. Mahecha,  
 652 P. Manning, T. Massad, B. E. Medlyn, J. Messier, A. T. Moles, S. C. Müller, K.  
 653 Nadrowski, S. Naeem, Ü. Niinemets, S. Nöllert, A. Nüske, R. Ogaya, J. Oleksyn,  
 654 V. G. Onipchenko, Y. Onoda, J. Ordoñez, G. Overbeck, W. A. Ozinga, S. Patiño,  
 655 S. Paula, J. G. Pausas, J. Peñuelas, O. L. Phillips, V. Pillar, H. Poorter, L. Poorter,  
 656 P. Poschlod, A. Prinzing, R. Proulx, A. Rammig, S. Reinsch, B. Reu, L. Sack, B.  
 657 Salgado-Negret, J. Sardans, S. Shiodera, B. Shipley, A. Siefert, E. Sosinski, J.-F.  
 658 Soussana, E. Swaine, N. Swenson, K. Thompson, P. Thornton, M. Waldram, E.  
 659 Weiher, M. White, S. White, S. J. Wright, B. Yguel, S. Zaehle, A. E. Zanne, and C.  
 660 Wirth. 2011. TRY—a global database of plant traits. *Global Change Biology* 17:  
 661 2905-2935.

662 Kazakou, E., D. Vile, B. Shipley, C. Gallet, and E. Garnier. 2006. Co-variations in litter  
 663 decomposition, leaf traits and plant growth in species from a Mediterranean old-  
 664 field succession. *Functional Ecology* 20: 21-30.

665 Kerkhoff, A. J., W. F. Fagan, J. J. Elser, and B. J. Enquist. 2006. Phylogenetic and  
 666 growth form variation in the scaling of nitrogen and phosphorus in the seed plants.  
 667 *American Naturalist* 168: E103-E122.

668 Koerselman, W., and A. F. M. Meuleman. 1996. The vegetation N: P ratio: a new tool

669 to detect the nature of nutrient limitation. *Journal of Applied Ecology* 33: 1441-  
670 1450.

671 Lambers, H., J. A. Raven, G. R. Shaver, and S. E. Smith. 2008. Plant nutrient-  
672 acquisition strategies change with soil age. *Trends in Ecology & Evolution* 23: 95-  
673 103.

674 Laughlin, D. C., P. Z. Fulé, D. W. Huffman, J. E. Crouse, and E. Laliberté. 2011.  
675 Climatic constraints on trait-based forest assembly. *Journal of Ecology* 99: 1489-  
676 1499.

677 Luo, Y. K. 2017. Functional traits of shrub species in eastern China. PhD thesis.  
678 University of Chinese Academy of Sciences. [In Chinese].

679 Mcgroddy, M. E., T. Daufresne, and L. O. Hedin. 2004. Scaling of C:N:P stoichiometry  
680 in forests worldwide: implications of terrestrial redfield-type ratios. *Ecology* 85:  
681 2390-2401.

682 Niinemets, Ü. 2001. Global-scale climatic controls of leaf dry mass per area, density,  
683 and thickness in trees and shrubs. *Ecology* 82: 453-469.

684 Niklas, K. J., T. Owens, P. B. Reich, and E. D. Cobb. 2005. Nitrogen/phosphorus leaf  
685 stoichiometry and the scaling of plant growth. *Ecology Letters* 8: 636-642.

686 Niklas, K. J. 2006. Plant allometry, leaf nitrogen and phosphorus stoichiometry, and  
687 interspecific trends in annual growth rates. *Annals of botany* 97: 155-163.

688 Ordonez, J. C., P. M. van Bodegom, J. P. M. Witte, R. P. Bartholomeus, J. R. van Hal,  
689 and R. Aerts. 2010. Plant strategies in relation to resource supply in mesic to wet  
690 Environments: Does Theory Mirror Nature? *American Naturalist* 175: 225-239.

691 Quested, H. M., J. H. C. Cornelissen, M. C. Press, T. V. Callaghan, R. Aerts, F. Trosien,  
692 P. Riemann, D. Gwynn-Jones, A. Kondratchuk, and S. E. Jonasson. 2003.  
693 Decomposition of sub-arctic plants with differing nitrogen economies: a functional  
694 role for hemiparasites. *Ecology* 84: 3209-3221.

695 Reich, P. B., and J. Oleksyn. 2004. Global patterns of plant leaf N and P in relation to



696 temperature and latitude. *Proceedings of the National Academy of Sciences of the*  
697 *United States of America* 101: 11001-11006.

698 Reich, P. B., J. Oleksyn, and I. J. Wright. 2009. Leaf phosphorus influences the  
699 photosynthesis-nitrogen relation: a cross-biome analysis of 314 species. *Oecologia*  
700 160: 207-212.

701 Reich, P. B., J. Oleksyn, I. J. Wright, K. J. Niklas, L. Hedin, and J. J. Elser. 2010.  
702 Evidence of a general 2/3-power law of scaling leaf nitrogen to phosphorus among  
703 major plant groups and biomes. *Proceedings of the Royal Society of London B:*  
704 *Biological Sciences* 277: 877-883.

705 Sardans, J., A. Rivas-Ubach, and J. Penuelas. 2012. The elemental stoichiometry of  
706 aquatic and terrestrial ecosystems and its relationships with organismic lifestyle  
707 and ecosystem structure and function: a review and perspectives. *Biogeochemistry*  
708 111: 1-39.

709 Sterner, R. W., and J. J. Elser. *Ecological stoichiometry: the biology of elements from*  
710 *molecules to the biosphere*. 2002; Press Princeton University Press, Princeton.

711 Stevens, P. F. Angiosperm Phylogeny Website. Version 14, July 2017 [and more or less  
712 continuously updated since]. <http://www.mobot.org/MOBOT/research/AP web />.

713 Sullivan, B. W., S. Alvarez-Clare, S. C. Castle, S. Porder, S. C. Reed, L. Schreeg, A. R.  
714 Townsend, and C. C. Cleveland. 2014. Assessing nutrient limitation in complex  
715 forested ecosystems: alternatives to large-scale fertilization experiments. *Ecology*  
716 95: 668-681.

717 Sundqvist, M. K., Z. F. Liu, R. Giesler, and D. A. Wardle. 2014. Plant and microbial  
718 responses to nitrogen and phosphorus addition across an elevational gradient in  
719 subarctic tundra. *Ecology* 95: 1819-1835.

720 Swaine, E. K. *Ecological and evolutionary drivers of plant community assembly in a*  
721 *Bornean rain forest*. PhD thesis, University of Aberdeen (2007).

722 Treseder, K. K. 2004. A meta-analysis of mycorrhizal responses to nitrogen, phosphorus,

723 and atmospheric CO<sub>2</sub> in field studies. *New Phytologist* 164: 347-355.

724 Tian, D., Z. B. Yan, K. J. Niklas, W. X. Han, J. Kattge, P. B. Reich, Y. K. Luo, Y. H.  
725 Chen, Z. Y. Tang, H. F. Hu, I. J. Wright, B. Schmid, and J. Y. Fang. 2018. Global  
726 leaf nitrogen and phosphorus stoichiometry and their scaling exponent. *National*  
727 *Science Review* 5: 728-739.

728 Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J.  
729 Cavender-Bares, T. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, E. Garnier,  
730 P. K. Groom, J. Gulias, K. Hikosaka, B. B. Lamont, T. Lee, W. Lee, C. Lusk, J. J.  
731 Midgley, M.-L. Navas, Ü. Niinemets, J. Oleksyn, N. Osada, H. Poorter, P. Poot, L.  
732 Prior, V. I. Pyankov, C. Roumet, S. C. Thomas, M. G. Tjoelker, E. J. Veneklaas,  
733 and R. Villar. 2004. The worldwide leaf economics spectrum. *Nature* 428: 821-  
734 827.

735 Wright, I. J., P. B. Reich, J. H. C. Cornelissen, D. S. Falster, E. Garnier, K. Hikosaka,  
736 B. B. Lamont, W. Lee, J. Oleksyn, N. Osada, H. Poorter, R. Villar, D. I. Warton,  
737 and M. Westoby. 2005. Assessing the generality of global leaf trait relationships.  
738 *New Phytologist* 166: 485-496.

739 Yan, Z. B., D. Tian, W. X. Han, Z. Y. Tang, and J. Y. Fang. 2017. An assessment on the  
740 uncertainty of the nitrogen to phosphorus ratio as a threshold for nutrient limitation  
741 in plants. *Annals of Botany* 120: 937-942.

742 Yang, X., X. L. Chi, C. J. Ji, H. Y. Liu, W. H. Ma, A. Mohhammat, Z. Y. Shi, X. P. Wang,  
743 S. L. Yu, M. Yue, and Z. Y. Tang. 2016. Variations of leaf N and P concentrations  
744 in shrubland biomes across northern China: phylogeny, climate and soil.  
745 *Biogeosciences* 13: 4429-4438.

746 Yang, X., W. M. Post, P. E. Thornton , and A. K. Jain. 2014. Global Gridded Soil  
747 Phosphorus Distribution Maps at 0.5-degree Resolution. ORNL DAAC, Oak  
748 Ridge, Tennessee, USA. 2014. <http://dx.doi.org/10.3334/ORNLDAAC/1223>.

749 Yang, Y. Z., H. Wang, S. P. Harrison, I. C. Prentice, I. J. Wright, C. H. Peng, and G. H.

750 Lin. 2019. Quantifying leaf-trait covariation and its controls across climates and  
751 biomes. *New Phytologist* 221: 155-168.

752 Zhang, S. B., J. L. Zhang, J. W. F. Slik, and K. F. Cao. 2012. Leaf element  
753 concentrations of terrestrial plants across China are influenced by taxonomy and  
754 the environment. *Global Ecology and Biogeography* 21: 809-818.